

INVESTIGATION AND MANAGEMENT OF GYNAECOMASTIA IN PRIMARY & SECONDARY CARE

PURPOSE

These guidelines are for the assessment, referral, and investigation of men with breast symptoms consistent with gynaecomastia, and cover the process from primary care to the specialist clinics of the Breast Unit and Endocrinology. They take into account the Best Practice Diagnostic Guidelines for Patients Presenting with Breast Symptoms from 2010.

BACKGROUND

Gynaecomastia is the enlargement of the male breast due to hyperplasia of the glandular tissue driven by alterations in male oestrogen:testosterone ratios. Pseudogynaecomastia is bilateral breast enlargement entirely due to adipose tissue. It does not require investigation or treatment.

Male breast cancer accounts for about 0.6% of all breast cancer: there almost 400 cases annually in the UK (c.f. 55,000 in women).

Benign gynaecomastia can be secondary to multiple medical and recreational drugs, as well as many chronic medical conditions.

- **Physiological**
 1. Neonatal: due to placental oestrogen transfer
 2. Pubertal: pubertal oestrogen production begins prior to testosterone production due to early maturation of aromatase (catalyzes conversion of androgens to oestrogens). Regression occurs in 90% of cases
 3. Senile: Age 70+. Up to 65% of men. Due to the reduction in testosterone relative to oestrogen
- **Drug induced** – 10-20% of gynaecomastia is due to prescribed drugs
 1. Oestrogen containing drugs eg. Bicalutamide, Buserelin, Goserelin
 2. Androgen receptor blocking drugs e.g. Cyproterone acetate, spironolactone, flutamide
 3. Androgen production inhibiting e.g. Finasteride, ketoconazole, dutasteride
 4. A list of medications that can cause gynaecomastia can be found in Appendix A
- **Drug induced** – recreational drugs such as marijuana, amphetamines, heroin, methadone
- **Pathological**
 1. Adrenal or testicular tumours <3% of gynaecomastia
 - a. Oestrogen or androgen producing tumours
 - b. Aromatase producing tumours
 - c. hCG producing tumours
 2. **Endocrine**
 - a. Primary hypogonadism [10% of gynaecomastia]
 - b. Secondary hypogonadism
 - c. Prolactinoma
 - d. Thyrotoxicosis
 - e. Acromegaly
 - f. Androgen insensitivity

SUMMARY STATEMENT: MANAGEMENT OF GYNAECOMASTIA IN PRIMARY & SECONDARY CARE

3. Systemic illness
 - a. Liver cirrhosis
 - b. Renal failure
 - c. Malnutrition
 - d. Obesity
 - e. HIV

INVESTIGATIONS RECOMMENDED TO BE DONE IN PRIMARY CARE

Before Referring

1. History to include:
 - Prescribed medications
 - Recreational drug use
 - Current and previous alcohol consumption
2. Chest wall examination – bilateral breasts

Do Not Investigate

- Adolescents with physiological pubertal gynaecomastia
- Elderly men with senile gynaecomastia
- Men with a drug related cause (prescribed medication or recreational drug use)
- Men with obvious breast cancer
- Men with fatty pseudogynaecomastia

Do Investigate

- Eccentric hard masses
- Rapid enlargement
- Recent onset in lean men >20 years
- Persistent painful gynaecomastia
- Massive gynaecomastia in adolescents
- Persistent gynaecomastia in adolescents, duration > 18-24 months

WHAT INVESTIGATIONS?

Blood tests

- 9am Testosterone, Thyroid Function Tests, Liver Function Tests, α -Fetoprotein, β -Human Chorionic Gonadotrophin
- If Testosterone is abnormal: Luteinizing Hormone, Follicle Stimulating Hormone, Sex Hormone Binding Globulin, albumin, oestradiol, prolactin
- Testicular Ultrasound Scan if any of the following abnormal blood results are noted: raised β HCG, raised α -Fetoprotein

GPS - WHEN AND WHERE TO REFER

Abnormal endocrine (hormonal) blood results

- Refer to Medical Endocrinology clinic

Abnormal β HCG or α FP blood results or abnormal finding on testicular USS

- Refer to Urology Clinic urgently

SUMMARY STATEMENT: MANAGEMENT OF GYNAECOMASTIA IN PRIMARY & SECONDARY CARE

Referral directly to the Breast Unit

In the presence of the following clinical scenarios, a referral directly to the local breast unit may be considered.

1. Clinical suspicion of malignancy
 - >50 year old man with unilateral firm sub-areolar mass with or without nipple discharge or with associated skin change
 - Bloody nipple discharge
 - Unilateral ulceration of the nipple
 - Urgent referral is appropriate
2. Unilateral lump with
 - No obvious physiological or drug cause
 - Increased risk - family history
 - Genetic conditions e.g. Klinefelter's Syndrome
3. Persistent painful gynaecomastia (>6 months) with normal blood tests

GYNAECOMASTIA IN THE BREAST UNIT

Gynaecomastia does not require all aspects of triple assessment

1. History:
 - Drug history
 - Alcohol history
 - Recreational drug use
 - Steroid use
 - Family history
2. Clinical examination:
 - Chest, bilateral
 - Nodal areas: axillae and supraclavicular fossae
 - Gynaecomastia can be described according to the Simon Classification (Appendix 1)
3. Imaging
 - Bilateral pseudogynaecomastia: No imaging
 - Bilateral gynaecomastia P2: No imaging
 - Unilateral lump in age <25years: No imaging
 - Unilateral lump in age >25 years *and* P2: No imaging
 - Unilateral lump in age >25years *and* P3+: USS +/- mammogram according to local practice
4. Pathology
 - Biopsy only if one or more of the following: P3+, M3+, U3+

HORMONAL TREATMENT

The patient must be informed that this treatment is off-licence. It is most effective for recent onset gynaecomastia, i.e. before gynaecomastia becomes fibrotic, and alleviates mastalgia, not always regression of the mass.

- Tamoxifen 10mg PO OD: 3-9 months.
- Anastrozole 1mg PO OD: 3 months.

SURGICAL REMOVAL


- Dependent on local CCG guidelines

Authors:	ABS Clinical Practice and Standards Committee
Produced:	March 2021
Version:	3

SUMMARY STATEMENT: MANAGEMENT OF GYNAECOMASTIA IN PRIMARY & SECONDARY CARE

APPENDIX 1

Simon classification for gynaecomastia



REFERENCES

Best Practice Diagnostic Guidelines for Patients Presenting with Breast Symptoms. Department of Health 2010

Braunstein GD. Gynaecomastia. *N Eng J Med.* 2007;357(12):1229-1237

Dall B et al. Overview of gynaecomastia in the modern era and the Leeds Gynaecomastia Investigation algorithm. *Breast J* 2011; 17(3):246-55

Gloucestershire Clinical Commissioning Group Policy for Male Breast Reduction for Gynaecomastia. 2015

Khan HN, Blamey RW. Endocrine treatment of physiological gynaecomastia. *BMJ* 2003;327:301-2

Mainiero MB, Lourenco AP, Barke LD, et al. ACR appropriateness criteria evaluation of the symptomatic male breast. *J Am Coll Radiol* 2015;12:678-82. doi:10.1016/j.jacr.2015

Newell MS et al. ACR Appropriateness Criteria Evaluation of the Symptomatic Male Breast. *J Am Coll Radiol.* 2015 Jul;12(7):678-82

Paredes M et al. Mammography and ultrasound in the evaluation of male breast disease. *Eur Radiol.* 2010;20(12):2797-2805

Perez EA et al. The role of mammography in male patients with breast symptoms. *Mayo Clin Proc.* 2007;82(3):297-300

Simon BE et al. Classification and surgical correction of gynaecomastia. *Plast Reconstr Surg.* 1973;51:48-52

Thiruchelvam P et al. Gynaecomastia – Clinical updates. *BMJ* 2016;354:i4833 doi:10.1136/bmj.i4833

Wallis M et al. The diagnostic value of clinical examination and imaging used as part of an age-related protocol when diagnosing male breast disease: an audit of 1141 cases from a single centre. *Breast* 2013 Jun;22(3):268-72

Zonderland HM et al. Overuse of imaging the male breast – findings in 557 patients. *Breast J* 2015 May-Jun;21(3):219-23

SUMMARY STATEMENT: MANAGEMENT OF GYNAECOMASTIA IN PRIMARY & SECONDARY CARE

APPENDIX 2

(taken from Thiruchelvam P, Walker JN, Rose K, Lewis J, Al-Mufti R. BMJ 2016;354:i4833)

DRUGS KNOWN TO CAUSE GYNAECOMASTIA	
Analgesics	opioid drugs (RA)
Antiandrogens	bicalutamide, flutamide, finasteride, dutasteride (AA)
Antifungals	ketoconazole (prolonged oral use) (AA)
Antihypertensives	calcium channel blockers (amlodipine, diltiazem, felodipine, nifedipine, verapamil) (UM), spironolactone (AA)
Antipsychotics (1st gen)	haloperidol (IP), olanzapine, paliperidone (high doses), risperidone (high doses), ziprasidone
Antiretrovirals	protease inhibitors (saquinavir, indinavir, nelfinavir, ritonavir, lopinavir), reverse transcriptase inhibitors (stavudine, zidovudine, lamivudine), efavirenz (UM)
Cardiovascular drugs	phytoestrogens (soya based products, high quantity) (E)
Chemotherapy drugs	methotrexate, alkylating agents—eg, cyclophosphamide, melphalan (AA); carmustine, etoposide, cytarabine, bleomycin, cisplatin (AA), vincristine (AA), procarbazine
Environmental exposures	phenothrin (antiparasitical)
Exogenous hormones	oestrogens (E), prednisone (male teenagers), human chorionic gonadotrophin (E), androgens (misuse by athletes) (E)
Gastrointestinal drugs	H ₂ histamine receptor blockers (cimetidine) (AA), proton pump inhibitors (eg, omeprazole) (AA)
Herbals	lavender, tea tree oil, dong quai (female ginseng), Tribulus terrestris, soy protein (300 mg/day), Urtica dioica (common nettle)
Recreational/illicit drugs	marijuana, amphetamines (UM), heroin (UM), methadone (UM), alcohol

DRUGS RARELY CAUSING GYNAECOMASTIA

Amiodarone (um)
 Aripiprazole, atorvastatin (um)
 Captopril (um), cetirizine, clonidine, cyproterone acetate (ishbg)
 Dasatinib, diazepam (ishbg), digoxin (e), domperidone, entecavir, fenofibrate (um)
 Fluoxetine (um)
 Gabapentin (aa)
 Imatinib (aa)
 Lisinopril, loratadine (aa)
 Metronidazole (aa), misoprostol (um)
 Paroxetine (um), penicillamine (aa), phthalates (um), pravastatin (um), pregabalin (aa)
 Ranitidine (aa), rosuvastatin (um)
 Sulindac, sulpiride, sunitinib (um)
 Theophylline (um)
 Venlafaxine (um)

SUMMARY STATEMENT: MANAGEMENT OF GYNAECOMASTIA IN PRIMARY & SECONDARY CARE

Codes	AA	Antiandrogenic
	RA	Reduced androgens
	E	Oestrogenic
	IAM	Increased androgen metabolism
	ISHBG	Increased concentration of sex hormone binding globulin
	IP	Increased prolactin
	UM	Unknown mechanism